



Cosyntropin (Cortrosyn)

Patient and Physician Information

Patient Name:	Date of Birth:	Patient Phone Number:
Physician Name:	Office Phone Number:	Fax Number:
Insurance:	Group Number:	Policy Number:
Hospitalization Status:	Patient Weight (kg):	Height (inches):
<input checked="" type="checkbox"/> Outpatient to Outpatient Infusion Center		
Allergies:		

Send patient demographics/insurance, clinical notes, and test results with orders

Diagnosis Code/Description for treatment: _____

Orders

Initiate IV Vascular Access Flush Orders #0643 for: ☐ Peripheral Line ☐ Midline ☐ PICC ☐ Port

☒ CORTISOL RESPONSE TO COSYNTROPIN x 3 –Baseline before injection, 30 minutes after injection, then 60 minutes after injection

☒ Cosyntropin (Cortrosyn) 0.25 MG INTRAVENOUS ONCE over 2 minutes.

Infusion Reaction

☒ If infusion reaction occurs, stop the infusion IMMEDIATELY, notify physician with details of reaction AND initiate the Outpatient Infusion HYPERsensitivity, OIC orders #1024

Discharge

☒ Discharge home 30 minutes after treatment complete if stable.

Date and Physician Signature

DATE: _____
11182507

TIME: _____

PHYSICIAN'S SIGNATURE