



## Immunoglobulin Intravenous (IVIG)

### Patient and Physician Information

<b>Patient Name:</b>	<b>Date of Birth:</b>	<b>Patient Phone Number:</b>
<b>Physician Name:</b>	<b>Office Phone Number:</b>	<b>Fax Number:</b>
<b>Insurance:</b>	<b>Group Number:</b>	<b>Policy Number:</b>
<b>Hospitalization Status:</b>	<b>Patient Weight (kg):</b>	<b>Height (inches):</b>
<input checked="" type="checkbox"/> Outpatient to Outpatient Infusion Center		
<b>Allergies:</b>		

\*\*\*Send patient demographics/insurance, clinical notes, and test results with orders\*\*\*

### Diagnosis Code/Description for treatment:

- ☐ Chronic Inflammatory Demyelinating Polyneuritis (G61.81)  
☐ Nonfamilial Hypogammaglobulinemia (D80.1) - Document IgG Level \_\_\_\_\_ mg/dL from what date: \_\_\_\_\_  
Document infection history (current/recurrent infection): \_\_\_\_\_  
☐ Immune Thrombocytopenic purpura (D69.3)  
☐ Other Primary Thrombocytopenia (D69.49)

### Orders

Initiate IV Vascular Access Flush Orders #0643 for: ☐ Peripheral Line ☐ Midline ☐ PICC ☐ Port

- ☒ Baseline vital signs then every 15 minutes x 4, then every 2 hours until infusion completed. If rate reduction is required, resume every 15 minute vital signs until symptoms resolved and vital signs returned to previous x 2.  
☒ Notify physician of: \*Heart rate 20% GREATER THAN baseline \*Respiratory distress or rate 20% GREATER THAN baseline  
\*Temperature GREATER THAN 38.5 degrees Celsius \* Systolic blood pressure 20% LESS THAN baseline \* Hives  
☒ Discontinue infusion and notify physician, if patient has: systolic blood pressure LESS THAN \_\_\_\_\_ mmHg  
☒ Normal Saline 0.9% Solution 20 milliliter/hour INTRAVENOUS (J7050 : 250 ML = 1 unit)

### Premedication

- ☐ Diphenhydramine (Benadryl) 25 MG ORAL ONCE 30 minutes prior to IVIG infusion (J1200 : 50 MG = 1 unit)  
☐ Acetaminophen (Tylenol) 325MG 2 TAB ORAL ONCE 30 minutes prior to IVIG infusion  
☐ methylPREDNISolone (Solu-Medrol) 125 MG IV PUSH ONCE 30 minutes prior to IVIG infusion. (J2930 : 125 MG = 1 unit)

### Infusion – Verify Insurance coverage to determine best product

- ☐ Privigen (J1459 : 500 MG = 1 unit) ☐ Octagam ((J1568 : 500 MG = 1 unit) ☐ Gammagard (J1569 : 500 MG = 1 unit)  
☒ Immune Globin (IgG) 10% \_\_\_\_\_ mg/kg (Patient's Ideal Body Weight) [TOTAL DOSE: \_\_\_\_\_ MG] at 30 milliliter/hour –Increase rate gradually by 30 mL/hr, allowing 15 to 30 minutes before each incremental increase. \*\* DO NOT EXCEED 150 mL/hr. If patient experiences flank pain, fever, tachycardia, bradycardia – REDUCE rate to the previous increment. (Dose will be rounded to the nearest vial size based on patient's current weight.)

### Infusion Reaction

- ☒ If infusion reaction occurs, stop the infusion IMMEDIATELY, notify physician with details of reaction AND initiate the Outpatient Infusion HYPERsensitivity, OIC orders #1024

### Discharge

- ☒ Discharge home 30 minutes after treatment complete if stable.

### Date and Physician Signature

DATE: \_\_\_\_\_  
07982508

TIME: \_\_\_\_\_

\_\_\_\_\_  
PHYSICIAN'S SIGNATURE