



## Methotrexate, Ectopic Pregnancy

### Patient and Physician Information

<b>Patient Name:</b>	<b>Date of Birth:</b>	<b>Patient Phone Number:</b>
<b>Physician Name:</b>	<b>Office Phone Number:</b>	<b>Fax Number:</b>
<b>Insurance:</b>	<b>Group Number:</b>	<b>Policy Number:</b>
<b>Hospitalization Status:</b>	<b>Patient Weight (kg):</b>	<b>Height (inches):</b>
<input checked="" type="checkbox"/> Outpatient to Outpatient Infusion Center		
<b>Allergies:</b>		

\*\*\*Send patient demographics/insurance, clinical notes, and test results with orders\*\*\*

### Diagnosis Code/Description for treatment:

- ☒ Ectopic Pregnancy

### Laboratory

- ☒ CBC WITH DIFFERENTIAL  
☒ COMPREHENSIVE METABOLIC PANEL  
☒ TYPE AND SCREEN if indicated to be RH negative  
☒ HCG, QUANTITATIVE  
☒ PROGESTERONE  
☒ RHIMMUNE GLOBULIN (Rhogam) prior to discharge, if indicated.

### Diagnostic Tests

- ☒ US ABDOMEN COMPLETE

### Orders

- ☒ Methotrexate 50 milligram/meter squared INTRAMUSCULAR ONCE x 1 dose (Dose: \_\_\_\_ MG = 50 x \_\_\_\_ BSA) – If total volume of the dose is GREATER THAN 2 milliliters, split the dose equally between 2 syringes (NOT to EXCEED 2 mL/syringe). Administer in EACH hip, MUST be administered by a nurse trained in administration and disposal of Methotrexate.  
**NOTE: DOSE is PER METER SQUARED, BASED on BODY SURFACE AREA (BSA)**

### Infusion Reaction

- ☒ If infusion reaction occurs, stop the infusion IMMEDIATELY, notify physician with details of reaction AND initiate the Outpatient Infusion HYPERsensitivity, OIC orders #1024

### Discharge

- ☒ Discharge home 30 minutes after treatment complete if stable.  
☒ Instruct patient to AVOID intake of Folic Acid including prenatal vitamins until advised by physician.  
☐ Schedule follow-up labs – CBC, Quantitative HCG, CMP in 4 days  
☐ Schedule follow-up labs – CBC, Quantitative HCG, CMP in 7 days

### Date and Physician Signature

DATE: \_\_\_\_\_  
08862508

TIME: \_\_\_\_\_

\_\_\_\_\_  
PHYSICIAN'S SIGNATURE