

Outpatient Infusion Center

Fax: 405-307-2244 Phone: 405-515-2470



Rituximab (Rituxan)

Patient and Physician Information		
Patient Name:	Date of Birth:	Patient Phone Number:
Physician Name:	Office Phone Number:	Fax Number:
Insurance:	Group Number:	Policy Number:
Hospitalization Status:	Patient Weight (kg):	Height (inches):
☑ Outpatient to Outpatient Infusion Center		
Allergies:		
Send patient demographics/insurance, clinical notes, and test results with orders		
Diagnosis Code/Description for treatment:		
Elaginosis Codo/Ecconplion for a caumona		
☐ Rheumatoid Arthritis, unspecified (M06.9)		
Orders		
Initiate IV Vascular Access Flush Orders #0643 for: ☐ Peripheral Line ☐ Midline ☐ PICC ☐ Port		
☑ Normal Saline 0.9% Solution 20 milliliter/hour INTRAVENOUS (J7050 : 250 ML = 1 unit)		
Other:		
Premedication Diphophydr MINE (Repedryl) 25 MC ORAL ONCE		
□ DiphenhydrAMINE (Benadryl) 25 MG ORAL ONCE□ DiphenhydrAMINE (Benadryl) 50 MG IV PUSH ONCE (J1200 : 50 MG = 1 unit)		
☐ Acetaminophen (Tylenol) 325MG 2 TAB ORAL ONCE ☐ methylPREDNISolone (Solu-Medrol) 125 MG IV PUSH ONCE – 30 minutes prior to infusion (J2930 : 125 MG = 1 unit)		
Interrity FIXE DIVISIONE (Sold-Media) 123 MG IV FOSTI ONCE - 30 minutes prior to midsion (32930 : 123 MG - 1 dilit)		
Infusion – Rituximab (Rituxan) [J9312 : 10 MG = 1 unit]		
☐ RITUximab (Rituxan) milligram/meter squared INTRAVENOUS over 3 to 8 hours. First dose – start at 50 MG/HR for 1 hour, then increase rate by 50 MG/HR EVERY 30 MINUTES. Subsequent doses – start at 100 MG/HR and increase by 100		
MG/HR. DO NOT EXCEED MAX rate of 400 MG/HR. Rheumatoid Arthritis		
☐ RITUximab (Rituxan) 1000 MG in 500 mL of 0.9% Normal Saline Solution (EXACT AMOUNT) INTRAVENOUS over 3 to 8		
hours. First dose – start at 50 MG/HR for 1		
doses – start at 100 MG/HR and increase by 100 MG/HR. Administer on day 1 and day 15. DO NOT EXCEED MAX rate of 400 MG/HR		
Infusion Reaction		
☑ If infusion reaction occurs, stop the infusion IMMEDIATELY, notify physician with details of reaction AND initiate the Outpatient		
Infusion HYPERsensitivity, OIC orders #1024 Discharge ☑ Discharge home 30 minutes after treatment complete if stable.		
Date and Physician Signature		
Date and I hydroidin dignature		
DATE: TIME:	_	PHYSICIAN'S SIGNATURE

Page 1 of 1

08142508